

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOLUTIONS OF EXERCISES.

ACKNOWLEDGMENTS.

Joseph Bowden, Jr. 325; H. Y. Benedict 326; Geo. R. Dean 322; W. H. Echols 322; A. Hall 327; J. E. Hendricks 325; Artemas Martin 314, 320; F. Morley 321; J. F. McCulloch 323; J. C. Nagle 326; W. B. Richards 282, 316, 322; W. O. Whitescarver 320; Chas. Yardley 320; De Volson Wood 324.

314*

A CYLINDER, diameter 2b, intersects a sphere, diameter 2a, the surface of the cylinder passing through the centre of the sphere. Required the part of the volume of the sphere contained by the cylinder.

[Artemas Martin].

SOLUTION.

Taking the origin at the centre of the sphere its rectangular equation is

$$x^2 + y^2 + z^2 = a^2, (1)$$

and that of the cylindric hole is

$$x^2 + y^2 = 2bx. \tag{2}$$

Also,

$$V = \iiint dx \ dy \ dz. \tag{3}$$

Let $x = r \cos \varphi$, and $y = r \sin \varphi$; whence (1) and (2) become

$$r^2 + z^2 = a^2, \tag{4}$$

$$r = 2b \cos \varphi,$$
 (5)

and

$$V = \iiint r \ dr \ d\varphi \ dz. \tag{6}$$

The limits of z are $-\sqrt{(a^2-r^2)}$ and $+\sqrt{(a^2-r^2)}$; of $r,2b\cos\varphi$ and 0; of $\varphi,\frac{1}{2}\pi$ and 0.

$$\begin{split} V &= 2 \int \int r \sqrt{(a^2 - r^2)} \, dr \, d\varphi, = -\frac{2}{3} \int (a^2 - r^2)^{\frac{3}{2}} \, d\varphi \\ &= \frac{4}{3} \int_0^{\frac{1}{2}\pi} a^3 d\varphi - \frac{4}{3} \int_0^{\frac{1}{2}\pi} (a^2 - 4b^2 \cos^2\!\varphi)^{\frac{3}{2}} \, d\varphi. \end{split} \tag{7}$$

^{*}The above solution is for the case a > 2b. The case a < 2b remains to be treated. The case a = 2b is old.—Ed.

134 solutions.

$$\begin{split} \text{Let } \varphi &= \frac{1}{2} \, \pi - \theta \, ; \text{ then } d\varphi = - \, d\theta, \cos \, \varphi = \sin \, \theta, \text{ and (7) becomes} \\ V &= \frac{2}{3} \, \pi a^3 - \frac{4}{3} \int \limits_0^{\frac{1}{2} \pi} (a^2 - 4b^2 \sin^2 \! \theta)^{\frac{3}{2}} \, d\theta \\ &= \frac{2}{3} \, \pi a^3 - \frac{4}{3} \, a^2 \int \limits_0^{\frac{1}{2} \pi} \sqrt{(a^2 - 4b^2 \sin^2 \! \theta)} \, d\theta \\ &+ \frac{16}{3} \, b^2 \int \limits_0^{\frac{1}{2} \pi} \sin^2 \! \theta \, \sqrt{(a^2 - 4b^2 \sin^2 \! \theta)} \, d\theta \end{split}$$

Putting e = 2b/a we have

$$V=rac{2}{3}\,a^3\pi-rac{16}{9}\,a\,(a^2-2b^2)~{
m E}\left[rac{1}{2}\,\pi,rac{2b}{a}
ight] \ -rac{4}{9}\,a\,(a^2-4b^2)~{
m F}\left[rac{1}{2}\,\pi,rac{2b}{a}
ight].$$
 When $b=rac{1}{2}\,a,~V=rac{2}{3}\,a^3igg[\pi-rac{4}{3}igg]$ [Artemas Martin].

A CIRCLE meets a hypocycloid of class 3 at six finite points. Show that the tangents to the hypocycloid at these six points touch a conic.

[Frank Morley.]

SOLUTION.

The equations of the hypocycloid in circular coordinates are

$$x/c = 2/t - t^2$$
, $y/c = 2t - 1/t^2$, (1)

t being a complex quantity of modulus 1. (See a paper on the Epicycloid, American Journal, Vol. XIII, No. 2).

Any circle is

$$xy + ax + \beta y + \gamma = 0.$$

solutions. 135

Substituting from (1), we have a sextic to determine t, and we observe that the product of the roots is 1.

The tangent to (1) at t is

$$ux + vy + 1 = 0,$$

where

$$u = -t/c (1 + t^3), \quad v = -t^2/c (1 + t^3).$$

Let the line equation of a conic be

$$(A, B, C, F, G, H)(u, v, 1)^2 = 0.$$

Substituting for u, v in terms of t, we have again

$$\overset{\scriptscriptstyle 6}{I\!\!I} t_r = 1,$$

 Π denoting a product. This condition ensures that the six tangents touch a conic; and we saw that it holds if the points t are concyclic.

[Frank Morley.]

322

The arc of a limaçon is shown in works on the Calculus to be equivalent to the arc of a certain ellipse. Show that the double point on the limaçon corresponds with Fagnani's point on the ellipse. [W. B. Richards.]

SOLUTION.

It is shown in works on the Calculus that

$$S = 2 \int\limits_0^{rac{1}{2}\pi} \{(a+b)^2\cos^2\!\!arphi + (a-b)^2\sin^2\!\!arphi\}^{rac{1}{2}}\,darphi,$$

is the quadrant of the ellipse on semi-axes 2(a + b) and 2(a - b), and also half of the whole length of the limaçon $r = a \cos \theta + b$.

It is well known that Fagnani's point divides the first arc into parts whose difference is 4b, while the half difference between the two loops of the limaçon is also 4b.

[W. H. Echols.]

Note.—George R. Dean also points out that at Fagnani's point $\cos \varphi = \sqrt{\frac{a+b}{2a}} = \cos \frac{1}{2}\theta$ at the node of the limacon.—Ed.

323

For solution see Analyst, Vol. I, No. 1, pp. 8-9. I proposed the problem in the Schoolday Visitor Magazine for May, 1872, nearly two years before Mr. Siverly used it in the Analyst. I had forgotten these facts when I sent it for publication in the Annals.

[Artemas Martin.]